

ORBITAL GRAPHS OF INFINITE PRIMITIVE PERMUTATION GROUPS

SIMON M. SMITH

ABSTRACT. If G is a group acting on a set Ω and $\alpha, \beta \in \Omega$, the digraph whose vertex set is Ω and whose arc set is the orbit $(\alpha, \beta)^G$ is called an *orbital digraph* of G . Each orbit of the stabiliser G_α acting on Ω is called a *suborbit* of G .

A digraph is *locally finite* if each vertex is adjacent to at most finitely many other vertices. A locally finite digraph Γ has more than one end if there exists a finite set of vertices X such that the induced digraph $\Gamma \setminus X$ contains at least two infinite connected components; if there exists such a set containing precisely one element, then Γ has *connectivity one*.

In this paper we show that if G is a primitive permutation group whose suborbits are all finite, possessing an orbital digraph with more than one end, then G has a primitive connectivity-one orbital digraph, and this digraph is essentially unique. Such digraphs resemble trees in many respects, and have been fully characterised in a previous paper by the author.

1. PRELIMINARIES

Throughout this paper, G will be a group of permutations of an infinite set Ω . A transitive group G is *primitive* on Ω if the only G -invariant equivalence relations admitted by Ω are the trivial and universal relations. A transitive group G is said to act *regularly* on Ω if $G_\alpha = 1$ for each $\alpha \in \Omega$.

If $\alpha \in \Omega$ and $g \in G$, we denote the image of α under g by α^g . Following this notation, all permutations will act on the right. The set of images of α under all elements of G is called an *orbit* of G ,

Date: February 2, 2008.

1991 *Mathematics Subject Classification.* 20 B 15; 05 C 25.

and is denoted by α^G . There is a natural action of G on the n -element subsets and n -tuples of Ω via $\{\alpha_1, \dots, \alpha_n\}^g := \{\alpha_1^g, \dots, \alpha_n^g\}$ and $(\alpha_1, \dots, \alpha_n)^g := (\alpha_1^g, \dots, \alpha_n^g)$ respectively.

If $\alpha \in \Omega$ we denote the stabiliser of α in G by G_α , and if $\Sigma \subseteq \Omega$ we denote the setwise and pointwise stabilisers of Σ in G by $G_{\{\Sigma\}}$ and $G_{(\Sigma)}$ respectively.

A *digraph* Γ will be a directed graph without multiple edges or loops; it is a pair $(V\Gamma, A\Gamma)$, where $V\Gamma$ is the set of *vertices* and $A\Gamma$ the set of *arcs* of Γ . The set $A\Gamma$ consists of ordered pairs of distinct elements of $V\Gamma$. Two vertices $\alpha, \beta \in V\Gamma$ are *adjacent* if either (α, β) or (β, α) lies in $A\Gamma$. A digraph is *locally-finite* if every vertex is adjacent to at most finitely many vertices. All paths will be undirected, unless otherwise stated. A digraph is *infinite* if its vertex set is infinite. The *distance* between two connected vertices α and β in Γ is denoted by $d_\Gamma(\alpha, \beta)$.

A *graph* is a pair $(V\Gamma, E\Gamma)$, in which $E\Gamma$ is a set of unordered pairs of elements of $V\Gamma$, each of which is known as an *edge* of Γ . The results presented in this paper concern digraphs, and are thus applicable to graphs.

The *automorphism group* of Γ will be denoted by $\text{Aut } \Gamma$. A graph or digraph is *primitive* if $\text{Aut } \Gamma$ is primitive on the set $V\Gamma$, and is *automorphism-regular* if $\text{Aut } \Gamma$ acts regularly on $V\Gamma$.

If G is a group of permutations of a set Ω and $\alpha, \beta \in \Omega$, then the digraph with vertex set Ω and arc set $(\alpha, \beta)^G$ is called an *orbital digraph* of G . Note such digraphs are necessarily arc-transitive. The orbital digraph $(\Omega, (\alpha, \beta)^G)$ is *non-diagonal* if α and β are distinct. Naturally, the graph $(\Omega, \{\alpha, \beta\}^G)$ is called an *orbital graph* of G .

The connected components of any digraph Γ are $\text{Aut } \Gamma$ -invariant equivalence classes on the vertex set of Γ ; thus any non-diagonal primitive orbital digraph is connected.

2. ENDS OF GRAPHS

A concept that is fundamental to the arguments contained herein is that of an *end*. A *half-line* L of a graph or digraph Γ is a one-way infinite cycle-free path $x_1x_2x_3\dots$ in Γ . The vertex x_1 is commonly

known as the *root* of L . There is a natural equivalence relation on the set of half-lines of Γ : two half-lines L_1 and L_2 are equivalent if there exists an infinite number of pairwise-disjoint paths connecting a vertex in L_1 to a vertex in L_2 . The equivalence classes of this relation are called the *ends* of Γ . An end is called *thick* if it contains infinitely many disjoint half-lines, otherwise it is called *thin*. By convention, a finite graph or digraph has no ends.

Theorem 2.1. ([10, Corollary 1.5]) *If Γ is an infinite locally finite connected vertex-transitive graph then Γ has 1, 2 or $2^{\aleph_0}$ ends.* $\square$

Given a digraph Γ , it is possible to construct an associated graph Γ' , whose vertex set is $V\Gamma$, with two vertices being adjacent in Γ' if and only if they are adjacent in Γ . Clearly, the ends of a digraph and its associated graph are the same. Thus, the above theorem is also true of digraphs.

Much of what is currently known about graphs with more than one end is underpinned by the work of Dicks and Dunwoody in their seminal text [2]. An accessible summary of their results pertaining to locally finite graphs can be found in [5].

The *connectivity* of an infinite connected graph or digraph Γ is the smallest possible cardinality of a subset W of $V\Gamma$ for which the induced digraph $\Gamma \setminus W$ is not connected. A *lobe* of Γ is a connected subgraph that is maximal subject to the condition it has connectivity strictly greater than one. If Γ has connectivity one, then the vertices α for which $\Gamma \setminus \{\alpha\}$ is not connected are called the *cut vertices* of Γ .

Let V_1 be the set of cut vertices of Γ , and let V_2 be a set in bijective correspondence with the set of lobes of Γ . Define T to be a bipartite digraph whose parts are V_1 and V_2 . Two vertices $\alpha \in V_1$ and $x \in V_2$ are adjacent in T if and only if α is contained in the lobe of Γ corresponding to x ; in this case there is a pair of arcs between them, one in each direction. This construction yields a tree, which is called the *block-cut-vertex tree* of Γ . If Γ has connectivity one and block-cut-vertex tree T , then for any group G acting on Γ , there is a natural action on T .

Two digraphs Γ_1 and Γ_2 are *quasi-isometric* if there exists a map $\phi : V\Gamma_1 \rightarrow V\Gamma_2$ and there exist constants $a, b > 0$ such that for all

$\alpha, \beta \in V\Gamma$ we have

$$\frac{1}{a} d_{\Gamma_1}(\alpha, \beta) - b \leq d_{\Gamma_2}(\phi(\alpha), \phi(\beta)) \leq a \cdot d_{\Gamma_1}(\alpha, \beta) + b.$$

In [4, Proposition 1], Möller shows if two locally finite connected graphs are quasi-isometric, then there is a natural bijection between their ends. This is also true of digraphs.

Lemma 2.2. *If G is a primitive group of permutations of an infinite set Ω and Γ_1 and Γ_2 are two non-diagonal orbital digraphs of G then there is a natural bijection between the ends of Γ_1 and the ends of Γ_2 .*

Proof. Suppose both $\Gamma_1 = (\Omega, (\alpha_1, \beta_1)^G)$ and $\Gamma_2 = (\Omega, (\alpha_2, \beta_2)^G)$ are locally finite orbital digraphs of G . Put $m_1 := d_{\Gamma_1}(\alpha_2, \beta_2)$ and $m_2 := d_{\Gamma_2}(\alpha_1, \beta_1)$, and let $a := \max\{m_1, m_2\}$. Since both Γ_1 and Γ_2 are connected, any arc in Γ_2 corresponds to a path of length m_1 in Γ_1 , and any arc in Γ_1 corresponds to a path of length m_2 in Γ_2 . Thus, if $\alpha, \beta \in \Omega$ then $d_{\Gamma_1}(\alpha, \beta) \leq m_1 \cdot d_{\Gamma_2}(\alpha, \beta) \leq a \cdot d_{\Gamma_2}(\alpha, \beta)$, and $d_{\Gamma_2}(\alpha, \beta) \leq m_2 \cdot d_{\Gamma_1}(\alpha, \beta) \leq a \cdot d_{\Gamma_1}(\alpha, \beta)$. Therefore $(1/a)d_{\Gamma_1}(\alpha, \beta) \leq d_{\Gamma_2}(\alpha, \beta) \leq a \cdot d_{\Gamma_1}(\alpha, \beta)$, so Γ_1 and Γ_2 are quasi-isometric. $\square$

Henceforth, no distinction will be made between an end of Γ_1 and its corresponding end in Γ_2 . Furthermore, we define a *permutation-end* of G to be an end of an orbital digraph of G . By the above lemma, the set of permutation-ends of G is equal to the set of ends of a non-diagonal orbital digraph of G , and is independent of the orbital digraph chosen.

3. THE CANONICAL ORBITAL DIGRAPH

In [6], Möller shows every connected locally finite primitive graph with more than one end is closely related to one with connectivity one.

Theorem 3.1. ([5, Theorem 15]) *If Γ is a connected locally finite primitive graph with more than one end then there exist vertices $\alpha, \beta \in V\Gamma$ such that the graph $(V\Gamma, \{\alpha, \beta\}^{\text{Aut } \Gamma})$ has connectivity one and each lobe has at most one end.* $\square$

Since any infinite vertex-transitive connectivity-one digraph has infinitely many ends, the following may be deduced.

Corollary 3.2. *If Γ is a locally finite primitive arc-transitive digraph then Γ has 0, 1 or $2^{\aleph_0}$ ends.* $\square$

Recall that the set of permutation-ends of a locally finite primitive group is equal to the set of ends of any one of its orbital digraphs. A consequence of the above corollary therefore, is that locally finite infinite primitive groups either have precisely one permutation-end, or they have uncountably many.

If G is a group of permutations of a set Ω , then the orbits of the action of the stabiliser G_α on Ω are called the α -suborbits of G . The cardinality of a suborbit is called a *subdegree* of G .

We can use Theorem 3.1 to tell us a great deal about the orbital digraphs of infinite primitive permutation groups.

Theorem 3.3. *If G is a primitive group of permutations of an infinite set Ω , with more than one permutation-end and no infinite subdegree, then G has a locally finite orbital digraph with connectivity one in which each lobe has at most one end.*

The proof of this theorem relies on the following five lemmas. Let G be a primitive group of permutations of an infinite set Ω .

Lemma 3.4. *If G has a locally finite orbital digraph with more than one end then G has an orbital digraph with connectivity one.*

Proof. Suppose G has a locally finite orbital digraph Γ with more than one end. Let Γ' be the graph associated with Γ . By Theorem 3.1, we may find vertices $\alpha, \beta \in \Omega$ such that the graph $(\Omega, \{\alpha, \beta\}^{\text{Aut } \Gamma'})$ has connectivity one. Now G is a primitive subgroup of $\text{Aut } \Gamma$, and thus $\text{Aut } \Gamma'$, so the graph $(\Omega, \{\alpha, \beta\}^G)$ is a connected subgraph of $(\Omega, \{\alpha, \beta\}^{\text{Aut } \Gamma'})$, and therefore has connectivity one. Hence, the digraph $\Gamma := (\Omega, (\alpha, \beta)^G)$ also has connectivity one. $\square$

Lemma 3.5. ([8, Lemma 2.1.]) *If G acts primitively on the vertices, and transitively on the arcs, of a connectivity-one digraph Γ , and Λ is any lobe of Γ with at least three vertices, then the subgroup of $\text{Aut } \Lambda$ induced by the setwise stabiliser $G_{\{\Lambda\}}$ acts primitively on the vertices of Λ .* $\square$

The following three lemmas are generalisations of several observations made in [6]. Although Möller restricts his attention to automorphism groups of graphs, for our purposes his arguments require only trivial modification.

Lemma 3.6. *If Γ is a locally finite connectivity-one orbital digraph of G and $\alpha, \beta \in V\Gamma$ lie in the same lobe Λ of Γ , then a lobe of $\Gamma' := (V\Lambda, (\alpha, \beta)^{G_{\{\Lambda\}}})$ is a lobe of $\Gamma' := (V\Gamma, (\alpha, \beta)^G)$. $\square$*

Let Γ be a locally finite orbital digraph of G with more than one end. Fix $\alpha, \beta \in V\Gamma$ such that $\Gamma^{(1)} := (V\Gamma, (\alpha, \beta)^G)$ has connectivity one. Let Λ_1 be a lobe of $\Gamma^{(1)}$ containing α , and let $G^{(1)} := G_{\{\Lambda_1\}}$. This group acts primitively on $V\Lambda_1$ by Lemma 3.5. If Λ_1 has more than one end then find $\beta_2 \in \Lambda_1$ such that $\Gamma^{(2)} := (V\Lambda_1, (\alpha, \beta_2)^{G^{(1)}})$ has connectivity one. Let Λ_2 be a lobe of $\Gamma^{(2)}$ containing α and put $G^{(2)} := G_{\{\Lambda_2\}}$. Again we note this group acts primitively on $V\Lambda_2$ by Lemma 3.5. For $i \geq 2$, if Λ_i has more than one end, find $\beta_{i+1} \in \Lambda_i$ such that $\Gamma^{(i+1)} := (V\Lambda_i, (\alpha, \beta_{i+1})^{G^{(i)}})$ has connectivity one. Let Λ_{i+1} be a lobe of $\Gamma^{(i+1)}$ containing α and put $G^{(i+1)} := G_{\{\Lambda_{i+1}\}}$. By Lemma 3.5, this group acts primitively on $V\Lambda_{i+1}$.

Lemma 3.7. *There is an $n \geq 1$ such that Λ_n has at most one end. $\square$*

Lemma 3.8. *For all integers $i \geq 1$, the digraph Λ_i is a lobe of the orbital digraph $(V\Gamma, (\alpha, \beta_i)^G)$. $\square$*

Proof of Theorem 3.3. Suppose G has more than one permutation-end and no infinite subdegree.

If Γ is a non-diagonal orbital digraph of G , then Γ has more than one end. Furthermore, if $(\alpha, \beta) \in A\Gamma$, and $(\gamma, \alpha) \in A\Gamma$, then α is adjacent to at most $|\beta^{G_\alpha}| + |\gamma^{G_\alpha}|$ vertices in Γ . Since G has no infinite subdegrees, this sum is finite, and since G acts transitively on $V\Gamma$, this digraph, and indeed all orbital digraphs of G , are locally finite.

Thus, by Lemma 3.4, G has a locally finite connectivity-one orbital digraph Γ . Using the notation described above, if we set $\Gamma^{(1)} := \Gamma$ then, by Lemma 3.7, there exists a positive integer n such that Λ_n has at most one end. Put $\Gamma' := (V\Gamma, (\alpha, \beta_n)^G)$. By Lemma 3.8, Λ_n is a lobe of Γ' , and since $\Lambda_n \neq \Gamma'$, we see Γ' has connectivity one. Furthermore,

G acts arc-transitively on Γ' , so every lobe of Γ' is isomorphic to Λ_n , and thus has at most one end. $\square$

The structure of primitive directed digraphs with connectivity one is well known.

Theorem 3.9. ([8, Theorem 2.6.]) *Let G be a vertex-transitive group of automorphisms of a connectivity-one digraph Γ whose lobes have at least three vertices. If G acts primitively on $V\Gamma$ and Λ is some lobe of Γ , then $G_{\{\Lambda\}}$ is primitive and not regular on $V\Lambda$.* $\square$

Recall that a digraph Γ is automorphism-regular if $\text{Aut } \Gamma$ acts regularly on $V\Gamma$.


Theorem 3.10. ([8, Theorem 3.3.]) *If Γ is a vertex-transitive digraph with connectivity one, then it is primitive if and only if the lobes of Γ are primitive but not automorphism-regular, pairwise isomorphic and each has at least three vertices.* $\square$

Combining Theorem 3.3, Theorem 3.9 and Theorem 3.10 we obtain the following.

Theorem 3.11. *If G is a primitive group of permutations of an infinite set Ω with more than one permutation-end and no infinite subdegree, then G has a locally finite orbital digraph Γ with connectivity one, whose lobes are primitive but not automorphism-regular, are pairwise isomorphic, have at least three vertices and at most one end. Furthermore, if Λ is a lobe of Γ , then $G_{\{\Lambda\}}$ acts primitively but not regularly on $V\Lambda$.* $\square$

If G is a primitive group of permutations of an infinite set Ω and Γ is a locally finite orbital digraph with connectivity one whose lobes have at most one end, then we shall call Γ a *canonical orbital digraph of G* . If Γ is such a digraph, and Λ is a lobe of Γ containing precisely one end ϵ , then ϵ will be called a *lobe-end* of Γ . If an end ϵ is not a lobe-end of Γ , it will be called a *tree-end*, as it corresponds to an end of the block-cut-vertex tree of Γ .

Observe that G permutes the lobe-ends of Γ transitively. Indeed, G acts arc-transitively on Γ , and each arc belongs to a unique lobe of Γ , so G permutes the lobes, and therefore the lobe-ends, transitively.

FIGURE 1. The trees T_1 and T_2

Since Γ has connectivity one, any thick ends of Γ must be lobe-ends. Thus, if Γ has a thick end then G acts transitively on the thick ends of Γ .

4. UNIQUENESS

Recall that if G is a primitive group and Γ_1 and Γ_2 are orbital digraphs of G then there is a natural bijection between the ends of Γ_1 and the ends of Γ_2 .

A *geodesic* between two connected vertices of a graph or digraph Γ is a path of minimal length between them. If T is a tree, and $\alpha, \beta \in VT$, there is exactly one geodesic in T between α and β , which we denote by $[\alpha, \beta]_T$. If ϵ is an end of T , then there is precisely one half-line in T that lies in ϵ with initial vertex α . We denote this half-line by $[\alpha, \epsilon)_T$. A connected subgraph C of Γ is said to *contain the end* ϵ if, for every half-line $(\alpha_i)_{i \in \mathbb{N}}$ lying in ϵ , we have $\alpha_i \in VC$ for all sufficiently large i .

Lemma 4.1. *Suppose Γ_1 and Γ_2 are canonical orbital digraphs of a primitive group G . If ϵ is a lobe-end of Γ_1 then ϵ is a lobe-end of Γ_2 .*

Proof. Let T_1 and T_2 be the lobe-cut-vertex trees of Γ_1 and Γ_2 respectively. Suppose, for a contradiction, that there is a lobe-end ϵ of Γ_1 that is a tree-end of Γ_2 . Since ϵ is a tree-end of Γ_2 , it is an end of T_2 . As it is a lobe-end of Γ_1 , there exists a lobe Λ_1 of Γ_1 whose end is ϵ , and a vertex $x \in VT_1 \setminus V\Gamma_1$ corresponding to the lobe Λ_1 .

The following argument is illustrated in Figure 1. Fix $\alpha \in V\Lambda_1$, and let C be the connected component of $T_2 \setminus \{\alpha\}$ containing ϵ . Since ϵ is an end of Λ_1 and ϵ is contained in C , there must be infinitely many vertices in $V\Lambda_1$ that lie in C . Choose $\alpha' \in V\Lambda_1 \setminus \{\alpha\}$ such that $\alpha' \in C$,

and fix any vertex $\delta \in [\alpha, \epsilon)_{T_2} \cap [\alpha', \epsilon)_{T_2}$. Let β be the vertex adjacent to x in $[x, \delta]_{T_1}$. Since β is adjacent to x in T_1 , we have $\beta \in V\Lambda_1$. If $\beta = \alpha$ then redefine $\alpha := \alpha'$. In this way, we can be sure $\beta \neq \alpha$.

Since $\beta \in V\Lambda_1$ and $G_{\{\Lambda_1\}}$ is primitive but not regular on $V\Lambda_1$, there is an element $g \in G_{\alpha, \{\Lambda_1\}}$ such that $\beta^g \neq \beta$. Now $\beta \in [\alpha, \delta]_{T_1}$, so $[\alpha, \delta]_{T_1} \neq [\alpha, \delta]_{T_1}^g$. However, $g \in G_\alpha$, so $[\alpha, \delta]_{T_1}^g = [\alpha, \delta^g]_{T_1}$ and therefore $\delta \neq \delta^g$. There is only one half-line in T_1 with initial vertex α contained in the end ϵ , so if $g \in G_{\alpha, \epsilon}$ then g must fix the half-line $[\alpha, \epsilon)_{T_1}$ pointwise. The vertex δ lies on this half-line, so $G_{\alpha, \epsilon} \leq G_\delta$. Since $\delta \neq \delta^g$, the automorphism g thus cannot lie in G_ϵ . This, however, is absurd, as ϵ is the end of Λ_1 , and $g \in G_{\alpha, \{\Lambda_1\}} \leq G_{\alpha, \epsilon}$. $\square$

If Γ is a locally finite connected graph or digraph, an automorphism $g \in \text{Aut } \Gamma$ is called *hyperbolic* if g fixes precisely two thin ends of Γ and leaves no non-empty finite subset of $V\Gamma$ invariant.

Theorem 4.2. ([9, Proposition 3.4]) *If T is an undirected tree and G is a group of automorphism of T containing no hyperbolic elements, then either G fixes some vertex or leaves some edge invariant, or G fixes some end of the tree T .* $\square$

Lemma 4.3. *Suppose G is a primitive group of permutations of an infinite set and Γ_1 and Γ_2 are canonical orbital digraphs of G . Let T_2 be the block-cut-vertex tree of Γ_2 . If Λ_1 is a lobe of Γ_1 then there exists $x \in VT_2$ such that $G_{\{\Lambda_1\}} \leq G_x$, and for all $\alpha_1, \alpha_2 \in V\Lambda_1$ we have $d_{T_2}(\alpha_1, x) = d_{T_2}(\alpha_2, x)$.*

Proof. If Λ_1 is infinite, then it contains exactly one lobe-end ϵ of Γ_1 , which by Lemma 4.1 is a lobe-end of Γ_2 . Let Λ_2 be the lobe of Γ_2 whose end is ϵ and let x be the vertex of T_2 corresponding to the lobe Λ_2 . Then $G_{\{\Lambda_1\}} = G_\epsilon = G_{\{\Lambda_2\}} = G_x$, so we have $G_{\{\Lambda_1\}}$ fixes some vertex in T_2 .

Now assume $V\Lambda_1$ is finite. Since $G_{\{\Lambda_1\}}$ cannot contain a hyperbolic element, we may use Theorem 4.2 to deduce that $G_{\{\Lambda_1\}}$ must fix a vertex of T_2 , or leave some edge of T_2 invariant, or fix an end of T_2 . In the latter two cases, the group must still fix some vertex of T_2 . Indeed, since no element of G may interchange two adjacent vertices of T_2 , if

the group $G_{\{\Lambda_1\}}$ leaves some edge $\{\alpha, x\}$ of T_2 invariant, then it must fix both α and x pointwise. Furthermore, if ϵ is some end of T_2 fixed by the group $G_{\{\Lambda_1\}}$, take x to be any vertex in T_2 such that the connected component of $T_2 \setminus \{x\}$ containing the end ϵ contains no element in $V\Lambda_1$. Since $V\Lambda_1$ is finite, there are infinitely many choices for such a vertex. Then any half-line in ϵ whose root lies in $V\Lambda_1$ must contain the vertex x . Thus, if $G_{\{\Lambda_1\}}$ fixes the end ϵ , it must also fix the vertex x .

Finally we observe that, since $G_{\{\Lambda_1\}}$ is transitive on $V\Lambda_1$, if $\alpha_1, \alpha_2 \in V\Lambda_1$ then $d_{T_2}(\alpha_1, x) = d_{T_2}(\alpha_2, x)$. $\square$

Henceforth, if Λ_1 is a lobe of Γ_1 , the vertex x in the above lemma will be referred to as the *centroid of Λ_1 in T_2* . Since the above arguments are symmetric in Γ_1 and Γ_2 , we can also make reference to the *centroid of Λ_2 in T_1* , where Λ_2 is any lobe of Γ_2 , and T_1 is the block-cut-vertex tree of Γ_1 .

Let Ω be some fixed infinite set and let G be a primitive group of permutations of Ω . Given any locally finite primitive connectivity-one orbital digraph Γ_1 of G , if each vertex lies in m distinct lobes and Λ is a lobe of Γ_1 , we write the digraph Γ_1 as $\Gamma(m, \Lambda)$. By Theorem 3.11, all locally finite primitive connectivity-one orbital digraphs of G can be written in this way. Since G acts transitively on the set of lobes of $\Gamma(m, \Lambda)$, there is a natural equivalence relation on the set of such digraphs: the orbital digraph Γ_2 is equivalent to Γ_1 if $\Gamma_2 = \Gamma(m, \Lambda')$ for some lobe Λ' of Γ_2 satisfying $V\Lambda = V\Lambda'$. It is under this equivalence relation that a canonical orbital digraph of a primitive permutation group can be considered essentially unique.

Theorem 4.4. *If G is a primitive group of permutations of an infinite set Ω possessing no infinite subdegree, then the canonical orbital digraphs of G are equivalent.*

Proof. The group G has at least one canonical orbital digraph by Theorem 3.11. Suppose G has two canonical orbital digraphs $\Gamma_1 = \Gamma(m_1, \Lambda_1)$ and $\Gamma_2 = \Gamma(m_2, \Lambda_2)$. By Theorem 3.10, both m_1 and m_2 are at least two, and the digraphs Λ_1 and Λ_2 are primitive but not automorphism-regular, and have at least three vertices. Let T_1 and T_2 be the block-cut-vertex trees of Γ_1 and Γ_2 respectively. We wish to show there is a lobe

Λ'_2 of Γ_2 such that $V\Lambda_1 = V\Lambda'_2$. From this we will deduce that the digraphs Γ_1 and Γ_2 are equivalent. Set $d := \sup\{d_{T_2}(\alpha, \beta) \mid \alpha, \beta \in V\Lambda_1\}$. This is finite, by Lemma 4.3, so we may fix $\alpha, \beta \in V\Lambda_1$ such that $d_{T_2}(\alpha, \beta) = d$. Let x_2 be the vertex of T_2 that is adjacent to α in the line $[\alpha, \beta]_{T_2}$. Since x_2 is adjacent to α in T_2 , it must lie in $VT_2 \setminus V\Gamma_2$, and therefore corresponds to some lobe Λ'_2 of Γ_2 . Let c_1 be the centroid of Λ_1 in T_2 .

We begin by showing $G_{\alpha, \{\Lambda_1\}} = G_{\alpha, \{\Lambda'_2\}}$. Now c_1 and β must lie in the same connected component of $T_2 \setminus \{\alpha\}$, and β and x_2 lie in the same component of $T_2 \setminus \{\alpha\}$, so c_1 and x_2 are in the same component of $T_2 \setminus \{\alpha\}$. Since x_2 is adjacent to α in $[\alpha, \beta]_{T_2}$, it follows that $x_2 \in [\alpha, c_1]_{T_2}$, and therefore $G_{\alpha, c_1} \leq G_{x_2}$. Thus

$$G_{\alpha, \{\Lambda_1\}} \leq G_{\alpha, c_1} \leq G_{\alpha, x_2} = G_{\alpha, \{\Lambda'_2\}}.$$

Now choose $\gamma \in \Lambda'_2 \setminus \{\alpha\}$ and repeat this argument, replacing the vertex β with γ , the tree T_2 with T_1 and the lobe Λ_1 with Λ'_2 . Whence, there exists a lobe Λ'_1 of Γ_1 containing α such that $G_{\alpha, \{\Lambda'_2\}} \leq G_{\alpha, \{\Lambda'_1\}}$, and therefore

$$G_{\alpha, \{\Lambda_1\}} \leq G_{\alpha, \{\Lambda'_2\}} \leq G_{\alpha, \{\Lambda'_1\}}.$$

Since G_α is transitive on the lobes of Γ_1 containing α , there exists $g \in G_\alpha$ such that $\Lambda_1^g = \Lambda'_1$. As there are only finitely many such lobes, $\Lambda_1^{g^n} = \Lambda_1$ for some natural number n . Now $G_{\alpha, \{\Lambda_1\}} \leq G_{\alpha, \{\Lambda_1\}^g}$, and therefore $G_{\alpha, \{\Lambda_1\}^g} \leq G_{\alpha, \{\Lambda_1\}^{g^2}}$. If we continue in this way we eventually obtain

$$G_{\alpha, \{\Lambda_1\}} \leq G_{\alpha, \{\Lambda_1\}^g} \leq \cdots \leq G_{\alpha, \{\Lambda_1\}^{g^n}};$$

however, $\Lambda_1^{g^n} = \Lambda_1$, so it follows that $G_{\alpha, \{\Lambda_1\}} = G_{\alpha, \{\Lambda_1\}^g}$. Thus

$$G_{\alpha, \{\Lambda_1\}} = G_{\alpha, \{\Lambda'_2\}}.$$

We now show $G_{\{\Lambda_1\}} = G_{\{\Lambda'_2\}}$. Let γ be the vertex in $[x_2, \beta]_{T_2}$ adjacent to x_2 , and observe $\gamma \in V\Lambda'_2$. Now $G_{\{\Lambda'_2\}}$ does not act regularly on $V\Lambda'_2$, so there is an element $g \in G_{\{\Lambda'_2\}}$ such that $\gamma^g \neq \gamma$. Therefore x_2 is the only vertex in $[x_2, \beta]_{T_2} \cap [x_2, \beta^g]_{T_2}$. Thus $d_{T_2}(\beta, \beta^g) = d_{T_2}(\beta, x_2) + d_{T_2}(\beta^g, x_2) = (d - 1) + (d - 1) = 2d - 2$. Now $g \in G_{\alpha, \{\Lambda'_2\}} = G_{\alpha, \{\Lambda_1\}}$, and $\beta \in V\Lambda_1$, so $\beta^g \in V\Lambda_1$. Furthermore, $d = d_{T_2}(\alpha, \beta)$ must be even, so if $d > 2$, then $d \geq 4$, and therefore $d_{T_2}(\beta, \beta^g) = 2d - 2 > d$. This is

not possible, as d was chosen to be maximal. It must therefore be the case that $d = 2$.

Since no two vertices of Λ_1 are at distance greater than d in T_2 , all vertices of Λ_1 are adjacent to x_2 in T_2 , and therefore lie in Λ'_2 . Now Λ_1 contains at least three vertices, each of which is adjacent to x_2 in T_2 , and so, since $G_{\{\Lambda_1\}}$ is transitive on $V\Lambda_1$, it must be the case that $G_{\{\Lambda_1\}}$ fixes the vertex x_2 , and therefore fixes $V\Lambda'_2$ setwise. Thus $G_{\{\Lambda_1\}} \leq G_{\{\Lambda'_2\}}$. If $G_{\{\Lambda_1\}} \neq G_{\{\Lambda'_2\}}$ then $G_{\alpha, \{\Lambda'_2\}} = G_{\alpha, \{\Lambda_1\}} < G_{\{\Lambda_1\}} < G_{\{\Lambda'_2\}}$, which cannot happen since $G_{\{\Lambda'_2\}}$ is primitive on $V\Lambda_2$. Hence $G_{\{\Lambda_1\}} = G_{\{\Lambda'_2\}}$.

Finally, since $G_{\{\Lambda_1\}}$ and $G_{\{\Lambda'_2\}}$ are transitive on $V\Lambda_1$ and $V\Lambda'_2$ respectively, and $\alpha \in V\Lambda_1 \cap V\Lambda'_2$, we have $V\Lambda'_2 = \alpha^{G_{\{\Lambda'_2\}}} = \alpha^{G_{\{\Lambda_1\}}} = V\Lambda_1$.

Now, Λ_1 is vertex-transitive, so the out-valency and in-valency of each vertex must be non-zero. Since Γ_1 is arc-transitive, this implies that G_α is transitive on the set of lobes of Γ_1 that contain α . Similarly, G_α is transitive on the set of lobes of Γ_2 containing α . Whence, $m_1 = |G_\alpha : G_{\alpha, \{\Lambda_1\}}| = |G_\alpha : G_{\alpha, \{\Lambda'_2\}}| = m_2$. Therefore $\Gamma_2 = \Gamma(m_1, \Lambda'_2)$ and $V\Lambda'_2 = V\Lambda_1$, so the digraphs Γ_1 and Γ_2 are equivalent. $\square$

It is now possible to describe every locally finite connectivity-one orbital digraph of a primitive group G .

Theorem 4.5. *If G is a primitive group of permutations of an infinite set Ω with more than one permutation-end and no infinite subdegree, then G has a canonical orbital digraph $\Gamma(m, \Lambda)$, and the canonical orbital digraphs of G are precisely the digraphs $\Gamma(m, \Lambda')$, where Λ' is an orbital digraph of $G_{\{\Lambda\}}$ acting on the set $V\Lambda$.*

Proof. Suppose G has a locally finite orbital digraph with more than one end. By Theorem 3.11, G has a canonical orbital digraph $\Gamma(m, \Lambda)$. Set $\Gamma := \Gamma(m, \Lambda)$, and fix vertices $\alpha, \beta \in V\Lambda$ such that $\Gamma = (\Omega, (\alpha, \beta)^G)$. Suppose Γ' is also a canonical orbital digraph of G . We show there exists an orbital digraph Λ' of $G_{\{\Lambda\}}$ acting on $V\Lambda$ such that Λ' is a lobe of Γ' and $\Gamma' = \Gamma(m, \Lambda')$. By Theorem 4.4, the digraph Γ' must be equivalent to Γ , so there exists a lobe Λ' of Γ' such that $\Gamma' = \Gamma(m, \Lambda')$ and $V\Lambda' = V\Lambda$. Let (α, β') be an arc in Λ' . Since Γ' is arc-transitive, $\Gamma' = (\Omega, (\alpha, \beta')^G)$. If (γ, δ) is any arc in Λ' , then there exists an element $g \in G$ such that $(\gamma, \delta) = (\alpha, \beta')^g$, but such an automorphism must fix

the lobe Λ' , and therefore lies in $G_{\{\Lambda'\}} = G_{\{\Lambda\}}$. Hence Λ' is an orbital digraph of $G_{\{\Lambda\}}$ acting on the set $V\Lambda$.

Conversely, suppose Λ' is an orbital digraph of $G_{\{\Lambda\}}$ acting on $V\Lambda$. We show the connectivity-one digraph $\Gamma(m, \Lambda')$ is a canonical orbital digraph of G .

It is simple to check that $G \leq \text{Aut } \Gamma(m, \Lambda')$, so we begin by showing G is arc-transitive on $\Gamma(m, \Lambda')$. Suppose Λ'_1 and Λ'_2 are lobes of $\Gamma(m, \Lambda')$. Then there exist lobes Λ_1 and Λ_2 of $\Gamma(m, \Lambda)$ such that $V\Lambda_1 = V\Lambda'_1$ and $V\Lambda_2 = V\Lambda'_2$. Since G acts arc-transitively on $\Gamma(m, \Lambda)$, it acts transitively on its lobes. Therefore, there exists an automorphism $g \in G$ such that $\Lambda_1^g = \Lambda_2^g$. Since $G \leq \text{Aut } \Gamma(m, \Lambda')$, we must also have $\Lambda_1'^g = \Lambda_2'$, so G permutes the lobes of $\Gamma(m, \Lambda')$ transitively. Thus G is transitive on the lobes of $\Gamma(m, \Lambda)$ and acts arc-transitively on each lobe, so it must act arc-transitively on the whole digraph $\Gamma(m, \Lambda')$.

It remains to show the lobes of $\Gamma(m, \Lambda')$ have at most one end. This follows immediately from Lemma 2.2, since both Λ and Λ' are orbital digraphs of $G_{\{\Lambda\}}$ acting on $V\Lambda$, they must have the same ends.

Hence, $\Gamma(m, \Lambda)$ is a connectivity-one orbital digraph of G in which each lobe has at most one end, and is therefore a canonical orbital digraph of the group G . $\square$

Thus, for any primitive group G possessing a canonical orbital digraph Γ , when speaking of the permutation-ends of G as being the ends of Γ , it makes sense to refer to the *lobe-ends* and *tree-ends* of G as being, respectively, the lobe-ends and tree-ends of its canonical orbital digraph. This can be extended to include any group G with just one permutation-end ϵ , by defining the *lobe-end* of G to be the end ϵ .

That a whole class of infinite primitive permutation groups possessing no infinite subdegree should each have an orbital digraph whose structure is tree-like is a property that will be explored and exploited in a future paper by the author. In particular, this result will be fundamental in the examination of the possible rates of subdegree growth of infinite primitive permutation groups.

The concluding section of this paper is conceived as a simple illustration of just one way in which the existence of the canonical orbital digraph may be used to easily obtain a deep understanding of the structure of infinite primitive groups possessing no infinite subdegree, through the application of Bass and Serre's theory on trees.

5. THE STRUCTURE OF INFINITE PRIMITIVE GROUPS

A common example of an amalgamated free product is when there are two groups G_1 and G_2 , with subgroups H_1 and H_2 that are isomorphic via $\varphi : H_1 \rightarrow H_2$. In this case the free product of G_1 and G_2 with amalgamated subgroup H_1 is denoted by $G_1 *_{H_1} G_2$.

Let G be a group acting on a digraph Γ . An inversion is a pair consisting of an element $g \in G$ and an arc $(x, y) \in A\Gamma$ such that $(x, y)^g = (y, x)$; if there is no such pair the group G is said to act *without inversion* on Γ . If G acts on Γ without inversion, we define the *quotient digraph* G/Γ to be the digraph whose vertex and arc set are the quotients of $V\Gamma$ and $A\Gamma$ respectively under the action of G ; a *fundamental domain* of $\Gamma \bmod G$ is a subgraph of Γ that is isomorphic to G/Γ . A *segment* of Γ is a subgraph of Γ consisting two adjacent vertices and an arc of Γ between them.

Theorem 5.1. ([7, Theorem 6]) *Let G be a group acting on a digraph Γ and let $P = \{(x, y), (y, x)\}$ be a segment of Γ . If P is a fundamental domain of $\Gamma \bmod G$ then Γ is a tree if and only if $G \cong G_x *_{G_{x,y}} G_y$. $\square$*

Using this powerful theorem, we may determine the structure of locally finite infinite primitive permutation groups with more than one permutation-end.

Theorem 5.2. *If G is a locally finite infinite primitive group of permutations of an infinite set Ω with more than one permutation-end, then G has a canonical orbital digraph $\Gamma(m, \Lambda)$ and*

$$G \cong G_\alpha *_{G_{\alpha, \{\Lambda\}}} G_{\{\Lambda\}},$$

where $\alpha \in V\Lambda$, and $G_{\alpha, \{\Lambda\}}$ is a non-trivial maximal proper subgroup of $G_{\{\Lambda\}}$ that fixes no element in $V\Lambda \setminus \{\alpha\}$.

Proof. Let G be an infinite primitive group possessing a finite suborbit whose pair is also finite, and suppose G has an orbital digraph with more than one end. By Theorem 3.11, G has a canonical orbital digraph Γ of the form $\Gamma(m, \Lambda)$, where $m \geq 2$ and Λ is a primitive but not automorphism-regular digraph, with at least three vertices and at most one end. Let T be the block-cut-vertex tree of Γ . Then $G \leq \text{Aut } T$. There is a natural bipartition of T , one part containing the vertices of Γ , and the other containing vertices corresponding to the lobes of Γ . Since G preserves this bipartition, the group G acts on T without inversion, and has two orbits on the vertices of T .

Let $x \in VT$ correspond with the lobe Λ of Γ , and fix $\alpha \in V\Lambda$. Then (α, x) is an arc in T . If we define $T' := (VT, (\alpha, x)^G)$, then T' is an arc-transitive tree upon which G acts without inversion, with fundamental domain the segment $\{\{\alpha, x\}, (\alpha, x)\}$. Hence, by Theorem 5.1,

$$G \cong G_\alpha *_{G_{\alpha, x}} G_x.$$

Since Λ is the lobe of Γ corresponding to the vertex x of T , the stabiliser G_x is equal to $G_{\{\Lambda\}}$. As $G_{\{\Lambda\}}$ acts primitively on $V\Lambda$, the group $G_{\alpha, \{\Lambda\}}$ is maximal in $G_{\{\Lambda\}}$. By Theorem 3.9, $G_{\alpha, \{\Lambda\}}$ fixes no vertex in $V\Lambda \setminus \{\alpha\}$. $\square$

This paper forms part of the author's DPhil thesis, completed at the University of Oxford, under the supervision of Peter Neumann. The author would like to thank Dr Neumann for his help and guidance. The author would also like to thank the EPSRC for generously funding this research.

REFERENCES

- [1] M. Bhattacharjee, D. Macpherson, R. G. Möller and P. M. Neumann, *Notes on infinite permutation groups*, Lecture Notes in Mathematics 1698 (Springer-Verlag, Berlin, 1998).
- [2] W. Dicks and M. J. Dunwoody, *Groups acting on graphs*, Cambridge Studies in Advanced Mathematics 17 (Cambridge University Press, Cambridge, 1989).
- [3] D. G. Higman, 'Intersection matrices for finite permutation groups', *J. Algebra* 6 (1967) 22–42.
- [4] R. G. Möller, 'Ends of graphs II', *Math. Proc. of the Cambridge Philos. Soc.* (3) 111 (1992) 455–460.

- [5] R. G. Möller, ‘Groups acting on locally finite graphs — a survey of the infinitely ended case’, *Groups ’93, Galway / St. Andrews, Vol. 2*, London Mathematical Society Lecture Note Series (Cambridge Univ. Press, Cambridge, 1993), pp. 426–456.
- [6] R. G. Möller, ‘Primitivity and ends of graphs’, *Combinatorica* (4) 14 (1994) 477–484.
- [7] J-P. Serre, *Trees* (Springer-Verlag, Berlin, 2003).
- [8] S. M. Smith, ‘Infinite primitive directed graphs’, *to appear*, <http://arxiv.org/abs/math.CO/0602011>.
- [9] J. Tits, ‘Sur le groupe des automorphismes d’un arbre’, *Essays on topology and related topics*, Mémoires dédiés à Georges de Rham (eds A. Haefliger and R. Narasimhan, Springer-Verlag, New York, 1970), pp. 188–211.
- [10] M. E. Watkins, ‘Ends and automorphisms of infinite graphs’, *Graph symmetry* (eds G. Hahn and G. Sabidussi), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. 497 (Kluwer Acad. Publ., Dordrecht, 1997), pp. 379–414.

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD

E-mail address: `simon.smith@chch.oxon.org`